

Most individuals who have already received a single dose of the bivalent vaccine are not currently eligible for another dose.

The FDA intends to make decisions about future vaccination after receiving recommendations on the fall strain composition at an FDA advisory committee in June.

Age	Received Monovalent Vaccine?	Received Bivalent Booster?	Which vaccine do they get?
65y+	Yes	Yes	2nd BIVALENT dose at least 4 months after last bivalent dose
6mo-64y*	Yes	Yes	No additional boosters at this time
12y+	Yes	No	1 dose of BIVALENT vaccine at least 2 months after last monovalent dose
12y+	No	No	1 dose of BIVALENT vaccine NOW
5y-11y	Yes	No	1 dose of BIVALENT vaccine at least 2 months after last monovalent dose
5y-11y	No	No	2 dose series of Moderna BIVALENT 6mo-5y vaccine 4-8 weeks apart OR 1 dose of Pfizer BIVALENT 5y-11y vaccine
6mo-4y	Yes – complete Pfizer series	No	1 dose of Pfizer BIVALENT 6mo-4y vaccine at least 2 months after series completion
6mo-5y	Yes- complete Moderna Series	No	1 dose of Moderna BIVALENT 6mo-5y vaccine at least 2 months after series completion
6mo-5y	Yes – incomplete series	No	Complete series with appropriate number of BIVALENT vaccines for age and vaccine brand
6mo-4y	No	No	3 dose series of Pfizer BIVALENT 6mo-4y vaccine; dose 1 and 2 given 3-8 weeks apart, dose 2 and 3 at least 8 weeks apart
6mo-5y	No	No	2 dose series of Moderna BIVALENT 6mo-5y vaccine 4-8 weeks apart
*Most immunocompromised individuals previously vaccinated and up to date with bivalent booster			2nd BIVALENT dose at least 2 months after last bivalent dose

- Most **unvaccinated individuals** may receive a single dose of a bivalent vaccine, rather than multiple doses of the original monovalent mRNA vaccines.
- **18+ yo:** Previously vaccinated with a monovalent COVID-19 vaccine who have not yet received a dose of a bivalent vaccine may receive a single dose of a bivalent vaccine.
- **65+ yo:** received a single dose of a bivalent vaccine may receive **one additional bivalent dose at least four months following their initial bivalent dose.**
- **Most individuals with certain kinds of immunocompromise who have received a bivalent COVID-19 vaccine** may receive 1 additional dose of a bivalent COVID-19 vaccine at least 2 months after a dose of a bivalent COVID-19 vaccine.
- **Children 6 months through 5 years of age who have received one, two or three doses of a monovalent COVID-19 vaccine** may receive a bivalent vaccine, but the number of doses that they receive will depend on the vaccine and their vaccination history (see below).
- **Unvaccinated Children:**
 - **6mo thru 4y:** may receive a 2-dose series of the Moderna bivalent vaccine (6 months through 5 years of age) **OR** a 3-dose series of the Pfizer-BioNTech bivalent vaccine (6 months through 4 years of age).
 - **5 year-olds:** may receive 2 doses of the Moderna bivalent vaccine **OR** only 1 dose of the Pfizer-BioNTech bivalent vaccine.